2024 UTME CHEMISTRY

OBJECTIVE STRUCTURED QUESTIONS

Which of the following produces effervescence with sodium hydrogen gen bonding D. ion-dipole forces carbonate? A. Ethanoic acid B. Ethane 17. The method suitable for separating fervescence with sodium hydrogen C. Bromine water D. Propene

What are the separation techniques involved in obtaining pure alum from powdered mixture of alum and sand? 18. The purity of common salt (NaCl) can A. Magnetic separation and centrifugation B. Dissolution, filtration and crystallization C. Distillation and chromatography D. Sedimentation and osmosis

What are the constituents of permalloy? A. Copper, zinc and tin B. Nickel and iron C. Copper, silver and manganese D. Copper, zinc and manganese

What is the colour obtained when starch iodide paper is introduced into a jar containing chlorine? A. Red B. 21. Blue C. Purple D. Blue-Black

The only factor that affects the internal energy of an ideal gas is A. pressure B. temperature C. volume D. number of moles

The element X that can combine with wgen to form an oxide of the form XO2 is A. sulphur B. fluorine C. chlone D. phosphorous

sust on the surface of a metal sheet mentains A. FeO B. hydrated Fe2O3 C.

radrate Fe₃O₄ D. FeCO₃

the food processing industry, the separation technique used for analyt- 22. ical purpose is A. crystallization B distillation C. chromatography D. centrifugation

An example of a solid-gas mixture is A. smoke B. foam C. fog D. mist

10. Which of the following will dissolve in water to give a solution with a pH greater than 7? A. NaCl B. Na2CO3 C. Na2SO4 D. NaHSO4

11. Which of the following metals react with water at the fastest rate? A. Calcium B. Magnesium C. Sodium D. Iron

12. The brown ring test is used to detect 24. the presence of A. nitrate ions B. sulphate ions C. chloride ions D. phosphate ions

13. The solubility of a solid in a liquid is increased by A. decreasing the tem- 25. perature B. increasing the pressure C increasing the surface area of the solid D. decreasing the volume of the sol-

The addition of charcoal to a filter bed in a town water supply is for the 26. purpose of A. removing heavy metals B. removing bacteria and viruses C. removing chlorine and organic compound D. softening the water.

15. The reaction between ethanol and phosphorous (V) chloride produces A. phosphorous (III) chloride B. phosphorous (V) oxide C. phosphorous 28. oxychloride D. phosphorous (III) ox-

The dominant intermolecular forces in ethanol are A. London dispersion forces B. Dipole-dipole forces C. hydro- 30. Which of the following is exothermic?

suspended particles in a liquid is A. distillation B. filtration C. magnetization D. Crystallization

be determined by A. measuring its
be determined by A. measuring its
A. 11.3 B. 2.7 C. 14 D. 3.54
boiling point B. measuring its density
32. Which of the following halogens is the C. measuring its melting point D. mea-

suring its solubility

19. The IUPAC name for the compound 33. In the electrolysis of dilute H₂SO₄, the HOCH₂CH₂CH₂OH is A. pentan-1 2-diol B. propan-1,3-diol C. propan-1,2-diol D. 1,2-dihydroxypropane 20. The purest form of natural water is A. 34. Which of the following gases can be

spring water B. river water C. lake wa-

ter D. rain water

- What is the main difference between 35. The main ore of aluminium is A. Gaempirical and molecular formulae? A. Empirical formula shows isotopes while molecular formula does not B. 36. Empirical formula shows the simplest whole-number ratio of atoms while molecular formula shows the actual number of atoms in a molecule. C. Empirical formulae are used for ionic compounds while molecular formulae 37. The number of unpaired electrons in a are used for covalent compounds D. Empirical formulae are used for gases while molecular formulae are used for 38. solids
- Which of the following bond types 39. involves the transfer of electrons? A. Hydrogen bonding B. Ionic bonding C. Covalent bonding D. Dipole-dipole interactions
- 23. librium constant (Kc) in a chemical reaction? A. It determines the type of catalyst used B. It determines the bond types in the reacting species C it determines the concentrations of the reactants and products D. None of the above

What is the term for the minimum amount of energy required for a chemical reaction to occur? A. Activation 1. energy B. Ionization energy C. Dissociation energy D. Kinetic energy

What is the term for the amount of heat energy required to raise the temperature of a unit mass of a substance 2. by 1 °C? A. Specific heat capacity B. Molar heat capacity C. Latent heat D. Heat of fusion.

The most suitable method for separat-3. ing a mixture of ethanol and water is A. filtration B. distillation C. sublima-

tion D. chromatography

27. The molecular formula of a compound 4. with an empirical formula CH2O and a molar mass of 180g/mol is A. C3H6O3 B. C₆H₁₂O₆ C. CH₄O D. C₄H₈O₂

The hybridization of carbon in ethene 5. is A. sp B. sp2 C. sp3 D. sp3d

Which of the following elements has the highest electronegativity? A. Sodium B. Fluorine C. Carbon D. Nitrogen

A. Melting of ice B. Dissolution of ammonium chloride in water C. Combustion of methane D. Evaporation of water

Determine the pOH of a solution of H₂SO₄ whose concentration is 0.001M

least reactive? A. Fluorine B. Iodine C. Chlorine D. Bromine

gas produced at the anode is A. hydrogen B. sulphur (IV) oxide C. oxygen D. hydrogen sulphide

collected over water? A. HCl B. NH;

C. CO₂ D. Cl₂

lena B. Bauxite C. Haematite D. Magnetite

Which of the following reactions is redox? A. NaOH + HCl → NaCl + H₂O B. $2Mg + O_2 \rightarrow 2MgO$ C. AgNO₃ + NaCl → AgCl + NaNO₃ D. $CaCO_3 \rightarrow CaO + CO_2$

ground-state nitrogen atom is A. 1 B.

2 C. 3 D. 4

The bond angle in methane is A. 90° B. 109.5° C. 120° D. 180°

Which of the following is not a property of transition metals? A. formation of coloured compounds B. High melting points C. Low density D. variable oxidation states

What is the significance of the equi- 40. The correct order of increasing boiling points for the following compounds is:

A. $CH_4 < C_2H_6 < C_3H_8 < C_4H_{10}$ B. C₄H₁₀<C₃H₈<C₂H₆<CH₄ C. C₂H₆<CH₄<C₄H₁₀<C₃H₈

D. C₃H₈<C₄H₁₀<CH₄<C₂H₆

2024 CHEMISTRY STRUCTURED QUESTIONS SOLUTIONS

A: Just like any other acid, ethanoic acid reacts with carbonate or hydrogen carbonate to give an effervescence of

CH3COOH+NaHCO3-CH3COONa+H2O+CO2 B: Alum is soluble in water while sand is not. So, dissolving the mixture in water, filtering and crystallizing can separate alum from sand

B: Permalloy is a nickel-iron magnetic alloy, composed of approximately 80% nickel and 20% iron. It is known for its high magnetic permeability

D: Starch iodide paper turns blueblack when exposed to chlorine gas. This is a common test for detecting the presence of chlorine gas.

B: The internal energy of an ideal gas is a function of its temperature only. This is a fundamental principle in thermodynamics.

6. A: Sulphur combines with oxygen to · form sulphur (IV) oxide (SO2), which is an acid anhydride. So, the element 19. B: X is carbon. Another element that fits this description is carbon, it forms CO2, another acid anhydride

7. B: When iron reacts with oxygen and 20. D: water, rust is formed. Rust is hydrated iron (III) oxide (Fe₂O₃. xH₂O)

C: Techniques like high-performance liquid chromatography (HPLC) and gas chromatography (GC) are commonly used to analyze food composition, detect contaminants and determine nutritional content.

A: Smoke is a solid-gas mixture, consisting of solid particles suspended in 21. B:

10. B:

- NaCl and Na2SO4 dissolve in water to produce a neutral solution (pH =

NaHSO₄ dissolves in water to produce an acidic solution (pH < 7)

Na₂CO₃ dissolves in water to pro- 22. B: duces an alkaline solution (pH > 7)

11. C: The reaction of sodium with water is really violent. It reacts with water in a highly exothermic reaction, releasing hydrogen gas and producing sodi- 23. um hydroxide solution.

The brown ring test is a chemical test used to detect the presence of nitrate ions (NO₃) in solution. It involves adding FeSO₄ and H₂SO₄ to the solu- 24. A: tion, which produces a brown ring if nitrate ions are present.

C: Increasing the surface area of a solid (probably by grinding) allows more particles to come into contact with the solvent, making it easier to dissolve

14. C: Activated charcoal is often added to filter beds in water treatment processes to remove chlorine, organic compounds and other impurities that can affect the taste and colour of the 26. B: water.

15. C:

The reaction between ethanol (C2H5OH) and phosphorous (V) chloride produces chloroethane (ethyl chloride), phosphorous oxychloride and hydrogen chloride. The balanced equation is:

 $C_2H_5OH+PCl_5\rightarrow C_2H_5Cl+POCl_3+HCl$

Ethanol is a polar molecule with a hydroxyl (-OH) group, which is capable of forming hydrogen bonds with other ethanol molecules. Hydrogen bonding is a strong intermolecular force.

17. B:

We can separate suspended particles from a liquid by passing the mixture 28. B: through a filter. The filter traps the suspended particles, allowing the clear liquid to pass through

18. C:

The purity of a solid can be determined by measuring its melting point. 29. B: A pure solid melts at a sharp, defined temperature or over a narrow tempera-

ture range while an impure solid melts 30. C: over a wide temperature range.

The IUPAC name of HOCH2CH2CH2OH is propan-1, 3-diol

Rain water is considered the purest form of water. It is formed through the earth's water cycle where water evaporates from the oceans, condenses into clouds and precipitates back to the earth as rain. This process distils the water, removing impurities and minerals, resulting in very low levels of contaminants

Empirical formula shows the simplest whole number ratio of atoms while molecular formula shows the actual number of atoms in a molecule. For instance, the molecular formula of benzene is C₆H₆ while its empirical 33. C: formula is CH.

Ionic/electrovalent bonding involves the transfer of electrons between at-

with opposite charges.

The equilibrium constant reflects the ratio of the concentrations of products to reactants at equilibrium. A high Kc value indicates that the products are present in greater proportion while a 35. B: low Kc value indicates otherwise

oms, resulting in the formation of ions

Activation energy is the minimum amount of energy required for a chem- 36. ical reaction to occur. It represents the energy barrier that must be overcome for the reaction to occur.

The specific heat capacity of a substance is the amount of heat energy required to raise the temperature of a 38. B: unit mass of a substance by 1 °C.

Ethanol and water can be separated by distillation because they have different boiling points. Ethanol boils at 78 °C while water boils at 100 °C.

With an empirical formula of CH₂O, the empirical formular mass

= 30g/mol.

To get the molecular formula, we multiply the ratio of the molar mass to the empirical formula mass by the empirical formula:

180gmol-1 $\frac{1}{30gmol^{-1}} = 6.$

Thus, the molecular formula $= (CH_2O)_6 = C_6H_{12}O_6.$

In ethene, each carbon atom forms three sigma bonds. There is also a Pi bond, leading to sp2 hybridization, where one p-orbital remains unhybridized.

Fluorine is the most electronegative element (of all known elements!)

All combustion processes are exothermic same is true for all neutralization reactions.

 $H_2SO_4 \rightarrow 2H^+ + SO_4^{2-}$ Then, $[H^+] = 2 \times 0.001M$ = 0.002 M $pH = -\log_{10}^{(0.002)} = 2.69897$

pH + pOH = 14pOH = 14 - 2.69897 = 11.3

32. B:

Reactivity of halogens decreases down the group due to the decreasing ability to attract electrons. Fluorine is the most reactive while iodine is the least reactive.

In the electrolysis of dilute H2SO4, oxygen is produced at the anode while hydrogen is produced at the cathode.

At the anode $4OH^- \rightarrow 2H_2O + O_2 + 4e^-$ At the cathode $2H^+ + 2e^- \rightarrow H_2$

CO2 is only slightly soluble in water and can therefore be collected over water, unlike highly soluble gases like HCl and NH₃.

Bauxite (Al₂O₃, nH₂O) is the primary ore from which aluminium is extracted (through electrolysis).

In the reaction, magnesium is oxidized and oxygen is reduced.

Methane has a tetrahedral geometry, where the bond angles are 109.5°. The hybridization of the carbon atom is sp³.

Transition metals are generally dense and have high melting points, they exhibit variable oxidation states and form coloured compounds.

As the number of carbon atoms increases in alkanes, the molecular mass increases, leading to stronger Van der Waals forces and higher boiling point